


POLAND


TOBACCO BURDEN FACTS

Poland ratified the Framework Convention on Tobacco Control on September 15, 2006.


Smoking prevalence in Poland (2010)¹¹



Tobacco use among youth (age 13-15) in Poland (2003)¹²



Deaths attributable to tobacco (2000)¹³



TOBACCO CONSUMPTION

Tobacco use is decreasing in Poland.

- A third (30%) of all adults (age 15+) smoke tobacco products with 37% of males and 24% of females smoking.¹
- Among youth (age 13-15), 19% smoke cigarettes (boys 20%; girls 17%).²
- 7% of youth (age 13-15) use tobacco products other than cigarettes (boys 9%; girls 5%).³
- In 2008, over 63 billion cigarettes were sold in Poland.⁴

HEALTH CONSEQUENCES

Tobacco use is deadly. Smoking kills up to half of all lifetime users.⁵

- Smoking kills approximately 70,000 people each year in Poland.⁶
- In 2000, 38% of male deaths occurring between the ages of 35-69 were attributed to smoking.⁷
- 90% of youth (age 13-15) are exposed to secondhand smoke in public places and 87% of youth are exposed to secondhand smoke in their homes.⁸

COSTS TO SOCIETY

Tobacco exacts a high cost on society.

- Healthcare costs associated with tobacco-related diseases amount to 18 billion PLN (6.4 billion USD) each year.⁹
- Losses in productivity and employment due to tobacco use are estimated to exceed 15 billion PLN (5.3 billion USD) each year.¹⁰
- Smoking also results in costs associated with fire damage and damage to the environment from the manufacturing and packaging of tobacco products.

1. Global adult tobacco survey (GATS): Poland. World Health Organization; 2009-2010. 2,3. CDC Global Youth Tobacco Survey (GYTS) [database on the Internet]. Poland - National 2003. Atlanta: Centers for Disease Control and Prevention. c2009 [cited 2009 August 17]. Available from: http://apps.nccd.cdc.gov/OSH_GTSS/default/Default.aspx. 4. Euromonitor International, (database online). London: Euromonitor; c2008 - (updated 2008 Sep 18; cited 2009 August 17). Industry reports: Tobacco- Poland. 5. Mackay J, Eriksen M, Shafey O. The Tobacco Atlas. 2nd ed. Atlanta: American Cancer Society; 2006. 6. Peto R, Lopez A, Boreham J, Thun M. Mortality from Smoking in Developed Countries 1950-2000. 2nd ed.; 2006. 7. Peto et al. 2006. 8. CDC GYTS, 2009. 9. Krzyżanowska A, Glogowski C. Nikotynizm na świecie: Następstwa ekonomiczne (The global nicotine addiction and its economic consequences). Menedżer Zdrowia; 2004 February. 98-103. 10. Krzyżanowska A, Glogowski C, 2004. 11. Nationwide survey on smoking behaviors and attitudes in Poland, 2002-2004 12. CDC GYTS, 2009. 13. Peto et al, 2006.